



NSW GP ALERT - whooping cough (pertussis)

Please distribute this information to all GPs and practice nurses

Whooping cough in school aged children means danger time for babies

3 key points for GPs

1. Whooping cough is increasing in NSW. Please consider the diagnosis in patients with cough.
2. People of any age can get whooping cough. Babies are at risk of severe disease.
3. Timely vaccination for babies is vital. Remember the first DTPa can be given at 6 weeks.

IDENTIFY POSSIBLE CASES

- Whooping cough has increased again in NSW, especially in school-aged children. Whooping cough immunity wanes over time and can occur in fully-vaccinated children.
- Cases classically present with bouts of coughing, followed by an inspiratory whoop and vomiting, but older children and adults may just have persistent bouts of coughing or a prolonged cough.
- Confirm the diagnosis by sending a nasopharyngeal swab for PCR. This is especially important for coughing adults and children who have contact with babies.

PROTECT OTHERS, ESPECIALLY BABIES

- Timely immunisation of infants is important because unvaccinated infants are at highest risk of infection and are also at high risk of complications.
- Because whooping cough immunity wanes over time, many older children and adults are susceptible and often transmit infections to infants.
- Free dTpa booster is still available for all new parents, grandparents and any other adults who will regularly care for infants less than 12 months of age.
- Advise all new parents to vaccinate their baby on time and keep their baby away from people with a cough.

PREVENT ITS SPREAD

- Whooping cough is easily spread within households, schools and childcare.
- Advise people with whooping cough to keep away from others and not attend school, childcare or work until no longer infectious. Cases are no longer infectious after 5 days of antibiotics or from 3 weeks after onset of symptoms.
- Recommended antibiotics: azithromycin, clarithromycin or erythromycin, see *Therapeutic Guidelines Antibiotic 2010*.
- Notify likely cases to your PHU. The PHU can help prevent further spread in high risk settings.

For more information, please call your local public health unit

Camperdown	02 9515 9420	Penrith	02 4734 2022	Broken Hill	08 8080 1499	Newcastle	02 4924 6477
Gosford	02 4349 4845	Randwick	02 9382 8333	Dubbo	02 6841 5569	Port Macquarie	02 6588 2750
Hornsby	02 9477 9400	Albury	02 6080 8900	Goulburn	02 4824 1837	Tamworth	02 6764 8000
Parramatta	02 9840 3603	Bathurst	02 6339 5601	Lismore	02 6620 7500	Wollongong	02 4221 6700

Pertussis factsheet for patients: www.health.nsw.gov.au/factsheets/infectious/pertussis.html

Dr Jeremy McAnulty
Director, Health Protection
14 October 2010

NSW Department of Health
ABN 92 697 899 630
73 Miller St North Sydney NSW 2060
Locked Mail Bag 961 North Sydney NSW 2059
Tel (02) 9391 9000 Fax (02) 9391 9101
Website www.health.nsw.gov.au